

Michael J. Ruiz, “Orthogonality Relation for Covariant Harmonic-Oscillator Wave Functions,” *Physical Review D* **10**, 4306 (1974).

The *American Physical Society (APS)* allows for the following. The “author or the author's employer may use all or part of the APS published article, including the APS-prepared version (e.g., the PDF from the online journal) without revision or modification, on the author's or employer's website as long as a fee is not charged. If a fee is charged, then APS permission must be sought. In all cases, the appropriate bibliographic citation and notice of the APS copyright must be included.” <https://journals.aps.org/copyrightFAQ.html>

“The author has the right to post and update the article on a free-access e-print server using files prepared and formatted by the author. Any such posting made or updated after acceptance of the article for publication by APS should include a link to the online APS journal article abstract. In all cases, the appropriate bibliographic citation and notice of the APS copyright must be included.” <https://journals.aps.org/copyrightFAQ.html>

The VOR (Version of Record) at the publisher’s site for this article is below.

<https://journals.aps.org/prd/abstract/10.1103/PhysRevD.10.4306>

Publications in APS journals are Copyright © American Physical Society.



*Work supported in part by the National Science Foundation under Grant No. GP-32998 X.

†On leave from the Institute for Theoretical Physics, University of Utrecht, The Netherlands.

‡Present address: Rockefeller University, New York, N. Y. 10021.

¹S. Weinberg, Phys. Rev. Lett. 29, 388 (1972); 29, 1698 (1972); Phys. Rev. D 7, 2887 (1973).

²I. Bars and K. Lane, Phys. Rev. D 8, 1169 (1973); 8, 1252 (1973).

³S. Y. Lee, J. M. Rawls, and L.-P. Yu, Nucl. Phys. B68, 255 (1974).

⁴J. Lieberman, Phys. Rev. D 9, 1749 (1974).

⁵B. de Wit, S.-Y. Pi, and J. Smith, this issue, Phys. Rev. D 10, 4278 (1974).

⁶P. W. Higgs, Phys. Lett. 12, 132 (1964); Phys. Rev. Lett. 13, 508 (1964); Phys. Rev. 145, 1156 (1966); F. Englert and R. Brout, Phys. Rev. Lett. 13, 321 (1964); G. S. Guralnik, C. R. Hagen, and T. W. B. Kibble, *ibid.* 13, 585 (1964); T. W. B. Kibble, Phys. Rev. 155, 1554 (1967).

⁷B. de Wit, Phys. Rev. D 9, 3399 (1974).

⁸H. Georgi and A. Pais, Phys. Rev. D 10, 1246 (1974).

Orthogonality relation for covariant harmonic-oscillator wave functions

Michael J. Ruiz

*Center for Theoretical Physics, Department of Physics and Astronomy,
University of Maryland, College Park, Maryland 20742*

(Received 20 May 1974)

Orthogonality relations for the Kim-Noz covariant harmonic-oscillator wave functions are discussed. It is shown that the wave functions belonging to different Lorentz frames satisfy an orthogonality relation. Furthermore, it is shown that for $n = m$ transitions there is a contraction factor of $(1 - \alpha^2)^{(n+1)/2}$, where α is the velocity difference between the two Lorentz frames.

The covariant harmonic-oscillator wave functions recently proposed by Kim and Noz¹ can be applied to a wide range of hadronic processes. The harmonic-oscillator characteristics prominently show up in hadronic mass spectra. The Lorentz-contraction properties of the oscillator wave functions can be seen in the nucleon elastic form factors and other electromagnetic transition amplitudes.²

In their paper, Kim and Noz are primarily concerned with the probability interpretation of their covariant wave functions. Their oscillator wave functions satisfy all the requirements of nonrelativistic quantum mechanics and enable us to extend the probability concept to the relativistic region. Kim and Noz, however, did not explicitly calculate the overlap integral of harmonic-oscillator wave functions belonging to two different Lorentz frames. The purpose of this note is to perform this overlap integral.

Kim and Noz start with the following differential equation:

$$\frac{1}{2} \left\{ -\nabla^2 + \frac{\partial^2}{\partial t^2} + \omega^2 [(\vec{x})^2 - t^2] \right\} \psi(x) = \lambda \psi(x). \quad (1)$$

This harmonic-oscillator equation is separable in the \vec{x} and t variables. Kim and Noz then observe that the above equation can also be written as

$$\frac{1}{2} \left\{ -\nabla_y^2 + \frac{\partial^2}{\partial y_0^2} + \omega^2 [(\vec{y})^2 - y_0^2] \right\} \psi(y) = \lambda \psi(y), \quad (2)$$

where the y variables are the Lorentz transforms of the x variables:

$$\begin{aligned} y_1 &= x_1, & y_2 &= x_2, \\ y_3 &= (1 - \beta^2)^{-1/2} (x_3 - \beta t), \\ y_0 &= (1 - \beta^2)^{-1/2} (t - \beta x_3). \end{aligned} \quad (3)$$

Equation (2) is also separable in the y variables. The normalizable solutions in the y variables are the Kim-Noz wave functions. Their wave function has the form

$$\begin{aligned} \psi_\lambda(y) &= N H_{n_1}(y_1) H_{n_2}(y_2) H_{n_3}(y_3) \\ &\quad \times \exp[-\frac{1}{2} \omega (\vec{y}^2 + y_0^2)], \end{aligned} \quad (4)$$

where $\lambda = \omega(n_1 + n_2 + n_3 + 1)$, and N is the normalization constant. The elimination of timelike oscillations can be done covariantly.¹ Since the transverse oscillations do not undergo Lorentz transformations, we shall assume that $n_1 = n_2 = 0$ in the following discussion.

The purpose of this note is to evaluate the following integral:

$$T_{nm}(\beta, \beta') = \int \psi_n(y) \psi_m(y') d^4x, \quad (5)$$

where y' variables are the y variables of Eq. (3) with β' . We can evaluate the above integral using the generating function for Hermite polynomials. This generating function has the form

$$\exp(-s^2 + 2sy_3) = \sum_{n=0}^{\infty} \frac{s^n}{n!} H_n(y_3). \quad (6)$$

We can now use the integral

$$I(s, r) = \left(\frac{\omega}{\pi}\right)^2 \int d^4x \exp(-s^2 + 2sy_3) \exp(-r^2 + 2ry'_3) \\ \times \exp\left[-\frac{1}{2}\omega(\vec{y}^2 + y_0^2 + \vec{y}'^2 + y_0'^2)\right] \quad (7)$$

to evaluate the integral in Eq. (5). Both y and y' are functions of \vec{x} and t . The transverse integrals can be performed trivially. For the t and z integrals we can use the variables ξ and η defined as

$$z = \frac{1}{\sqrt{2}}(\xi + \eta), \quad t = \frac{1}{\sqrt{2}}(\xi - \eta) \quad (8)$$

to complete the square of the exponent of the Gaussian factor. We obtain

$$I(s, r) = (1 - \alpha^2)^{1/2} \exp[2rs(1 - \alpha^2)^{1/2}], \quad (9)$$

where $\alpha = (\beta - \beta')(1 - \beta\beta')^{-1}$. Since the power-series expansion of Eq. (9) contains only equal powers of r and s , the integral of Eq. (5) vanishes unless $m = n$. When $m = n$, we find from the coefficient of $(rs)^n$ in the expansion of Eq. (9)

$$T_{nm}(\beta, \beta') = (1 - \alpha^2)^{(n+1)/2}. \quad (10)$$

The above results can be expressed in the following orthogonality relation³:

$$T_{nm}(\beta, \beta') = (1 - \alpha^2)^{(n+1)/2} \delta_{nm}. \quad (11)$$

The above result suggests that the covariant harmonic oscillators behave like nonrelativistic oscillators if they are in the same Lorentz frame. If two oscillators are in different frames, the orthogonality is preserved. The ground-state oscillator wave function, consisting of one half-wave, is contracted by $(1 - \alpha^2)^{1/2}$. The n th excited state, consisting of $(n + 1)$ half-waves, is contracted by $(1 - \alpha^2)^{(n+1)/2}$.

The author would like to thank Professor Y. S. Kim for suggesting this problem.

¹Y. S. Kim and M. E. Noz, Phys. Rev. D 8, 3521 (1973).

²R. Lipes, Phys. Rev. D 5, 2849 (1972). Lipes calculates his transition amplitudes in the Lorentz frame where the excited-state resonance is at rest, and his wave functions coincide with those of Kim and Noz in this particular frame. For this reason, the use of the

Kim-Noz wave function will lead to Lipes's calculation.

³The first attempt to get the Lorentz contraction factor was made by Markov, who obtained a similar result for $n = 0$ and $\beta' = 0$. See M. Markov, Nuovo Cimento Suppl. 3, 760 (1956).